

Non-Abelian Statistics
in the Fractional Quantum Hall States *

X. G. Wen

School of Natural Sciences
Institute of Advanced Study
Princeton, NJ 08540

ABSTRACT: The Fractional Quantum Hall states with non-Abelian statistics are studied. Those states are shown to be characterized by non-Abelian topological orders and are identified with some of the Jain's states. The gapless edge states are found to be described by non-Abelian Kac-Moody algebras. It is argued that the topological orders and the associated properties are robust against any kinds of small perturbations.

PACS numbers: 05.30.-d, 73.20.Dx

* Research supported by DOE grant DE-FG02-90ER40542
Published in *Phys. Rev. Lett.* **66**, 802 (1991)

It becomes more and more clear that the ground states of strongly interacting electron systems may contain very rich structures^{1,2,3,4,5,6} which cannot be characterized by broken symmetries and are called the topological orders.² Physical characterizations of the topological orders are discussed in Ref. 2,6. It is shown that the Fractional Quantum Hall (FQH) states, the chiral spin states and anyon superfluid states contain non-trivial topological orders characterized by the Abelian Chern-Simons (CS) theories.^{2,3,5,7} It is interesting to know whether the non-Abelian (NA) topological orders characterized by the NA CS theories⁸ can be realized in strongly interacting electron systems or not. In this paper we will construct some FQH states which contain NA topological orders. The effective theory of these states is shown by NA CS theory and the quasiparticles carry NA statistics⁸ We will also discuss how to understand the NA statistics in terms of the electron wave function.

Different electron wave functions with filling fraction $1/2n$ have been constructed in Ref. 4. The quasiparticles in these states were shown to have NA statistics, provided that these states are incompressible and the quasiparticles have finite size for a local Hamiltonian.

The spirit of our discussion is very similar to that in the mean field approach of the spin liquid states.^{9,10} The similar construction is also used to study the $SU(N)$ spin chains.¹¹ Consider a two dimensional spinless (*i.e.*, spin polarized) electron system in strong magnetic field with filling fraction $\nu = M/N$. For convenience we will put the electron system on a lattice, thus the electron Hamiltonian has a form

$$H = \sum_{ij} [t_{ij} e^{ieA_{ij}} c_i^\dagger c_j + V_{ij} n_i n_j] \quad (1)$$

where A_{ij} is the electromagnetic gauge potential on the lattice and $n_i = c_i^\dagger c_i$. To construct a FQH state with a NA topological order, we would like to break each electron into N partons¹² ψ_a each carrying electric charge e/N :

$$c = \psi_1 \psi_2 \dots \psi_N = \frac{1}{N!} \sum_{ab\dots c} \epsilon_{ab\dots c} \psi_a \psi_b \dots \psi_c \quad (2)$$

where ψ_a are fermionic fields and N is odd. After substituting (2) into (1) and making a mean field approximation we reach the following mean field Hamiltonian

$$H_{\text{mean}} = \sum_{ij} t_{ij} e^{ieA_{ij}/N} U_{ij,ab} \psi_{ia}^\dagger \psi_{jb} \quad (3)$$

where

$$U_{ij,aa'} = e^{ieA_{ij} \frac{N-1}{N}} \left(\frac{1}{N!} \right)^2 \langle \epsilon_{ab\dots c} (\psi_b \dots \psi_c)^\dagger \epsilon_{a'b'\dots c'} (\psi_{b'} \dots \psi_{c'})_j \rangle \quad (4)$$

The mean field solution U_{ij} can be obtained by minimizing the average of the Hamiltonian (1) on the ground state of H_{mean} in (3) (*i.e.*, $E = \langle \Phi_{\text{mean}} | H | \Phi_{\text{mean}} \rangle$). Let us assume that there exists a Hamiltonian H such that the mean field solution takes the most symmetric form $U_{ij,ab} = \eta \delta_{ab}$. In this case the mean field Hamiltonian (3) describes N kinds of free partons in magnetic field, each with a filling fraction $\nu = M$. Thus the mean field ground

state wave function is given by $\Phi_{\text{mean}}(z_i^a) = \prod_{a=1}^N \chi_M(z_i^a)$ where z_i^a is the coordinate of the a -th kind of the partons and $\chi_M(z_i)$ is the fermion wave function of M filled Landau levels.

Notice that the mean field theory (3) contains a lot of unphysical degrees of freedom arisen from the breaking of the electrons into partons. In order to use the mean field theory to describe the original electron system we need to project into the physical Hilbert space which satisfies the constraint

$$\psi_{1i}^\dagger \psi_{1i} = \dots = \psi_{Ni}^\dagger \psi_{Ni} \quad (5)$$

In the physical Hilbert space, different kinds of the partons always move together. The bound states of the partons correspond to the original electrons. The electron ground state wave function can be obtained by doing the projection on the mean field wave function Φ_{mean} by setting $z_i^1 = z_i^2 = \dots = z_i$ where z_i is the electron coordinate. The electron wave function obtained this way is just one of the FQH wave functions propose by Jain.¹² In the following we will call such a state NAF (NA FQH) state.

The wave function of the NAF state, $[\chi_M(z_i)]^N$, is the exact ground state of the following local Hamiltonian.¹³ The kinetic energy in the Hamiltonian is such that the first $NM - N + 1$ Landau levels have zero energy and other Landau levels have finite positive energies. One such kinetic energy is given by $\prod_{I=0}^{NM-N} [K - (I + \frac{1}{2})\omega_c]$ where $K = -\frac{1}{2m}(\partial_i - ieA_i)^2$. The two-body potential in the Hamiltonian has a form $V(r) \propto \partial^{2N-2}\delta(r)$. One can easily see that the Hamiltonian is positive definite, and the NAF state has zero energy because the electrons in the ground state all lie in the first $NM - N + 1$ Landau levels and the ground state wave function has N -th order zeros as $z_i \rightarrow z_j$. However it is not clear whether the state has the *highest* filling fraction among the zero energy states (this is related to the incompressibility). We can only show that among the Jain's states¹² the NAF state is the zero energy state with highest filling fraction. We do not know whether it is sufficient to only consider the Jain's states. It would be interesting to numerically test the incompressibility of the NAF state for the above Hamiltonian. Numerical calculations has only been done for the projection into first two Landau levels.¹⁴ In this case one indeed find the Jain's 2/5 state to be the exact *incompressible* ground state.

The projection, or the constraint (5), can be realized by including a gauge field. Notice that under local $SU(N)$ transformations $\psi_{ai} \rightarrow W_{i,ab}\psi_{bi}$, $W_i \in SU(N)$, the electron operator c_i in (2) is invariant. Thus the Hamiltonian contain a *local* $SU(N)$ symmetry after we substitute (2) into (1). The local $SU(N)$ symmetry manifest itself as a gauge symmetry in the mean field Hamiltonian (3). Notice that (3) is invariant under the $SU(N)$ gauge transformation $W_i: \psi_i \rightarrow W_i\psi_i$ and $U_{ij} \rightarrow W_i U_{ij} W_j^\dagger$. The gauge fluctuation in the mean field theory can be included by replacing the mean field value $U_{ij} = \eta$ by $U_{ij} = \eta \exp(ia_{ij})$ where a_{ij} is a $N \times N$ hermitian matrix. a_{ij} is just the $SU(N)$ gauge potential on the lattice. The time component of the $SU(N)$ gauge field can be included by adding a term¹⁰ $\psi_i^\dagger a_0(i)\psi_i$ to the mean field Hamiltonian. The constraint (5) is equivalent to the following constraint¹⁰

$$J_\mu^I(i) = 0, \quad I = 1, \dots, N^2 - 1 \quad (6)$$

where J_μ^I are the $SU(N)$ charge and the current density. The constraint (6) can be enforced in the mean field theory by integrating out the gauge field fluctuation a_μ .¹⁰ After

the projection, the only surviving states are those which are invariant under the local $SU(N)$ transformations. Those states correspond to the physical electron states.

The effective theory of the NAF state described above can be obtained by first integrating out ψ_a field:

$$\mathcal{L}_{0\text{eff}} = \frac{M}{4\pi N} A_\mu \partial_\nu A_\lambda \epsilon^{\mu\nu\lambda} + \frac{M}{8\pi} \text{Tr} a_\mu f_{\nu\lambda} \epsilon^{\mu\nu\lambda} \quad (7)$$

which is just the level M $SU(N)$ CS theory.⁸ $f_{\mu\nu}$ in (7) is the strength of the $SU(N)$ gauge field. The quasiparticle excitations in the NAF state correspond to the holes in various Landau levels of the partons. Those excitations are created by the parton fields ψ_a . After including the gauge fields, the properties of the quasiparticles are described by the following effective Lagrangian

$$\mathcal{L}_{q\text{eff}} = \sum \psi^\dagger [(i\partial_t + \frac{e}{N} A_0 + a_0) - \frac{1}{2m} (\partial_i - i\frac{e}{N} A_i - ia_i)^2] \psi \quad (8)$$

(7) and (8) describe the low energy properties of the NAF state.

The NA CS theory given by (7) and (8) have been studied in detail in Ref. 8. The quasiparticles ψ_a (which are called the Wilson lines in Ref. 8) are found to have NA statistics. In the following we will summarize some special properties associated with the NA statistics and discuss their relation to the microscopic electron wave function. Let us put the NAF state on a sphere. The ground state of (7) is found to be non-degenerated on the sphere. (On genus g Riemann surface the ground states are degenerate.) Now let us create m quasiparticles and m' quasiholes using the operators ψ_{a_i} and $\psi_{a_j}^\dagger$. If we have ignored the gauge field a_μ (setting $a_\mu = 0$), the Hilbert space generated by $\psi_{a_i}|_{i=1}^m$ and $\psi_{a_j}^\dagger|_{j=1}^{m'}$ would be $(\mathcal{H}_R)^m \times (\mathcal{H}_{\bar{R}})^{m'}$ which has $N^{m+m'}$ dimensions. Here \mathcal{H}_R is the fundamental representation of the $SU(N)$ and $\mathcal{H}_{\bar{R}}$ is the dual of \mathcal{H}_R . However after we include the gauge fluctuations and do the projection $z_i^a = z_i$, only the gauge invariant states can survive the projection and appear as the physical states of the original electron system. In particular all the states that transform non-trivially under the global $SU(N)$ are project away. Thus the Hilbert space $\mathcal{H}_{mm'}$ of the physical states is contained in the $SU(N)$ invariant subspace of $(\mathcal{H}_R)^m \times (\mathcal{H}_{\bar{R}})^{m'}$.⁸ $\text{Inv}((\mathcal{H}_R)^m \times (\mathcal{H}_{\bar{R}})^{m'})$ In the above we have only used the global $SU(N)$ gauge symmetry. The local gauge symmetry may further reduce the dimension of the Hilbert space. Not every (global) $SU(N)$ singlet state can survive the projection and become a physical state. Thus the dimension of $\mathcal{H}_{mm'}$ can be less than that of $\text{Inv}((\mathcal{H}_R)^m \times (\mathcal{H}_{\bar{R}})^{m'})$.

When $m = 1$ and $m' = 0$, there is no invariant state and the dimension of \mathcal{H}_{10} is zero. When $m = m' = 1$ there is only one invariant state. It is shown that such a state is always physical and \mathcal{H}_{11} is one dimensional. In this case moving one particle around the other induces a Berry's phase $\exp(\frac{i2\pi(N+1)}{N(N+M)})$. When $m = m' = 2$, $\text{Inv}((\mathcal{H}_R)^2 \times (\mathcal{H}_{\bar{R}})^2)$ is two dimensional. It turns out that \mathcal{H}_{22} is two dimensional if $M > 1$ and one dimensional if $M = 1$.⁸ As we interchange the two particles created by ψ_{a_i} , $i = 1, 2$, we obtain a NA Berry's phase for $M > 1$. The 2×2 matrix describing the NA Berry's phase is found⁸ to have eigenvalues $-\exp(\frac{i\pi(-N+1)}{N(N+M)})$ and $\exp(\frac{i\pi(N+1)}{N(N+M)})$. For $M = 1$ the Hilbert space \mathcal{H}_{22} is one dimensional and the corresponding Berry's phase is $\exp(i\pi/N)$. The later result

is expected because the $M = 1$ NAF state is just the Laughlin state with filling fraction $1/N$. The reproduction of the well known results of the Laughlin states is a non-trivial self consistency check of our theory.

Some of the above results can be easily understood in terms of the microscopic electron wave function. First we notice that the mean field state Φ_{mean} is a (global) $SU(N)$ singlet and the NAF wave function can be expressed as $\langle 0 | \prod_i c(z_i) | \Phi_{\text{mean}} \rangle = [\chi_M(z_i)]^N$ where $c(z_i)$ is given by (2). The quasiparticles discussed above are described by the following electron wave function $\langle 0 | \prod_i c(z_i) \prod_{I,I'} \psi_{a_I} \psi_{b_{I'}}^\dagger | \Phi_{\text{mean}} \rangle$. Since $\langle 0 | \prod_i c(z_i)$ is an $SU(N)$ singlet, it is clear that only the states in $\text{Inv}((\mathcal{H}_R)^m \times (\mathcal{H}_{\bar{R}})^{m'})$ can survive the projection and give rise to non-zero electron wave functions. The dimension of the physical Hilbert space may be smaller than that of the invariant space because the electron wave functions induced from different mean field singlet states may not be orthogonal to each other. For $m = m' = 1$ the electron wave function can be obtained by the projection of the mean field state $\psi_1(Z_1)\psi_1^\dagger(Z_2)|\Phi_{\text{mean}}\rangle$. The electron wave function is non-zero and is given by $\chi_M(z_i; Z_1; Z_2)[\chi_M(z_i)]^{N-1}$, where $\chi_M(z_i; Z_1; Z_2)$ has one hole and one particle at Z_1 and Z_2 . Thus \mathcal{H}_{11} is one dimensional. For $m = m' = 2$ the two electron wave functions $\Phi_{1,2}$ can be obtained by the projection of mean field states $\psi_1(1)\psi_1(2)\psi_1^\dagger(3)\psi_1^\dagger(4)|\Phi_{\text{mean}}\rangle$ and $\psi_1(1)\psi_2(2)\psi_1^\dagger(3)\psi_2^\dagger(4)|\Phi_{\text{mean}}\rangle$ (which contain the two singlets). Notice that locally the electron wave function are the same near each quasiparticle no matter the quasiparticle is created by ψ_1 or ψ_2 . More precisely the physical correlation functions, like the density correlation, are the same around each quasiparticle when the quasiparticles are well separated. This is because ψ_1 can be rotated into ψ_2 by a global $SU(N)$ transformation while the density correlation being a $SU(N)$ invariant quantity will not be changed. The effects of the other quasiparticles can be ignored since the correlation in the NAF states is short ranged and the other particles are far away. Thus the two electron states Φ_1 and Φ_2 should have the same local correlations and hence the same energy. Such a degeneracy is a bulk property just like the degeneracy of the FQH states on a torus.

When $M = 1$ each kind of partons only fill the first Landau level. The action of $\psi_1(1)\psi_1(2)\psi_1^\dagger(3)\psi_1^\dagger(4)$ on the first-Landau-level wave function corresponds to multiplication of a factor $A_{22} = \sum_{i_1, i_2} \delta(Z_3 - z_{i_1})\delta(Z_4 - z_{i_2}) \prod_{i, j \neq i_1, i_2} \frac{(z_i - Z_1)(z_j - Z_2)}{(z_i - z_{i_1})(z_j - z_{i_2})}$ and the action of $\psi_1(1)\psi_1^\dagger(3)$ corresponds to a factor $A_{11} = \sum_{i_1} \delta(Z_3 - z_{i_1}) \prod_{i \neq i_1} \frac{z_i - Z_1}{z_i - z_{i_1}}$. After the projection the two resulting electron wave functions are given by $A_{22}(\chi_M)^N$ and $(A_{11})^2(\chi_M)^N$ which describe the same state since $A_{22} \propto (A_{11})^2$. Similar derivations apply to other values of m and m' and the physical Hilbert space $\mathcal{H}_{mm'}$ can be shown to be at most one dimensional for $M = 1$. This is just the result of the NA CS theory. More detailed discussions of the structures of excitations in the NAF state will appear elsewhere.

We would like to remark that although the gauge field mediate no long range interactions between quasiparticles due to the CS term, the quasiparticles ψ_a are not really equivalent to the “free” quarks in absence of the gauge field. This is because the quasiparticles are dressed by NA flux which carries the $SU(N)$ charge. Thus it is conceivable that when $M = 1$ the quasiparticles behave like Abelian anyons with no internal degree of freedom, as has been shown in the above discussion.

Now let us discuss another fascinating property of the NAF state – the gapless edge excitations^{15,6} in the NAF state. We will follow the discussions in Ref. 6. First let us

ignore the constraint (6) and set $a_\mu = 0$ in the mean field theory. In this case the edge excitations are those of the IQH states¹⁵ described by

$$\mathcal{L} = \sum_{a\alpha} i\lambda^{a\alpha\dagger}(\partial_0 - v\partial_x)\lambda^{a\alpha} \quad (9)$$

where $\lambda^{a\alpha}$ is a fermion field describing the edge excitations of the α -th Landau level of the a -th kind of the partons. The Hilbert space of (9) can be represented¹¹ as a direct product of the Hilbert spaces of a $U(1)$ Kac-Moody (KM) algebra,¹⁶ a level N $SU(M)$ KM algebra and a level M $SU(N)$ KM algebra. This decomposition is a generalization of the spin-charge separation in the 1D Hubbard model. Notice that the total central charge of the above three KM algebras is $1 + \frac{N(M^2-1)}{M+N} + \frac{M(N^2-1)}{N+M} = MN$ which is equal to the central charge of (9). The above three KM algebra are generated by currents $J_\mu = eN^{-1}\lambda^{a\alpha\dagger}\partial_\mu\lambda^{a\alpha}$ (the electric current), $j_\mu^I = t_{\alpha\beta}^I\lambda^{a\alpha\dagger}\partial_\mu\lambda^{a\beta}$, and $J_\mu^I = T_{ab}^I\lambda^{a\alpha\dagger}\partial_\mu\lambda^{b\alpha}$, where t^I (T^I) are the generators of the $SU(M)$ ($SU(N)$) Lie algebra. The currents in the $SU(N)$ KM algebra are just the currents in (6) which couple to the $SU(N)$ gauge field a_μ . To obtain the physical edge excitations in the electron wave function, we need to do the projection to enforce the constraint (6). Because of the above decomposition, the projection can be easily done by removing from the Hilbert space of (9) the states associated with the $SU(N)$ KM algebra.⁶ The remaining physical edge states are generated by the $U(1) \times SU(M)$ KM algebra. The central charge of the $U(1) \times SU(M)$ KM algebra is given by $c = M\frac{MN+1}{M+N}$ and the specific heat (per unit length) of the edge excitations¹⁷ is $C = c\frac{\pi T}{6v}$. The electron creation operator⁶ on the edge is given by $c = \lambda^{1\alpha_1} \dots \lambda^{N\alpha_N}$ which has a propagator $(x - vt)^{-N}$ along the edge. We would like to point out that in general the edge excitations may have several different velocities in contrast to what was implicitly assumed above.

The above construction can be easily generalized in a number of directions: A) We may decompose electrons into partons with different electric charge. B) We may choose a different mean field ground state which break the $SU(N)$ gauge symmetry. Actually A) is a special case of B).¹⁸ The effective CS theory for A) and B) will in general contain several Abelian and NA gauge fields. In particular the FQH states studied in Ref. 3,6 correspond to breaking the $SU(N)$ gauge symmetry into $[U(1)]^{N-1}$ gauge symmetry. One interesting NAF state in the case A) is the $\nu = \frac{1}{1+\frac{1}{2}+\frac{1}{2}} = \frac{1}{2}$ state. Its NA statistics is described by the level 2 $SU(2)$ CS theory. The electrons in such a state lie in the first three Landau levels.

We would like to argue that the NAF states studied in this paper are generic states and their NA topological orders are robust against small perturbations. A) The NA structures in the NAF states come from the $SU(N)$ gauge symmetry of the mean field ground state. To destroy the NA topological orders (and the associated NA statistics) we must break the $SU(N)$ gauge symmetry through the Higgs mechanism. This cannot be achieved unless we add *finite* perturbations. B) All excitations in the NAF have finite energy gap and the interactions between them have finite range. Therefore the NAF states do not have infrared divergences and it is self consistent to assume the interactions between the excitations do not destabilize the NAF states. Thus we expect the properties studied in this paper are universal properties of the NAF states which are robust against any small perturbations. The NAF states are a new type of the infrared fixed points and the NA topological orders should appear as a general possibility for the ordering in the ground states of strongly interacting electron systems.

It is not clear under what condition the NAF states might be realized in nature. However since the NAF states are generic states, they may appear in experiments under right conditions, especially when the electron density is low and higher Landau levels are important. The low density FQH states are largely unexplored in experiments.

REFERENCES

1. F.D.M. Haldane *Phys. Rev. Lett.* **51**, 605 (1983); B. Halperin, *Phys. Rev. Lett.* **52**, 1583 (1984).
2. X.G. Wen, *Int. J. Mod. Phys. B* **2**, 239 (1990); X.G. Wen, *Phys. Rev.* **B40**, 7387 (1989); X.G. Wen and Q. Niu, *Phys. Rev.* **B41**, 9377 (1990).
3. B. Blok and X.G. Wen, *Phys. Rev.* **B42**, 8133 (1990); *Phys. Rev.* **B42**, 8145 (1990); IASSNS-HEP-90/66, IAS preprint.
4. G. Moore and N. Read, (unpublished).
5. N. Read, *Phys. Rev. Lett.* **65**, 1502 (1990)
6. X.G. Wen, “edge excitations of the FQH states with generic filling fractions”, IAS preprint IASSNS-HEP-90/42.
7. X.G. Wen and A. Zee, *Nucl. Phys.* **B15**, 135 (1990).
8. E. Witten, *Comm. Math. Phys.* **121**, 351 (1989).
9. X.G. Wen, F. Wilczek and A. Zee, *Phys. Rev.* **B39**, 11413 (1989). R. Laughlin and Z. Zou, *Phys. Rev.* **B41**, 664 (1990).
10. I. Affleck, Z. Zou, T. Hsu and P.W. Anderson, *Phys. Rev. B* **38**, 745 (1988); E. Dagotto, E. Fradkin and A. Moreo, *Phys. Rev. B* **38**, 2926 (1988).
11. I. Affleck, *Nucl. Phys.* **B265**, 409 (1986).
12. J.K. Jain, *Phys. Rev.* **B40**, 8079 (1989).
13. A similar observations has also been made by J.K. Jain (private communication).
14. Private communication from A.H. MacDonald.
15. B.I. Halperin, *Phys. Rev.* **B25**, 2185 (1982). X.G. Wen, “Gapless boundary excitations in the quantum Hall states and in the chiral spin states”, NSF-ITP-89-157, ITP preprint (Oct. 1989); A.H. MacDonald, *Phys. Rev. Lett.* **64**, 220 (1990); X.G. Wen, *Phys. Rev. Lett.* **64**, 2206 (1990); X.G. Wen, *Phys. Rev.* **B41**, 12838 (1990); M. Stone, “Edge waves in the QH effect”, IL-TH-8, Urbana preprint (1990); M. Stone, “Vertex Operators in the QH effects”, IL-TH-90-32, Urbana preprint (1990); F.D.M. Haldane, *Bulletin of APS* **35**, 254 (1990).
16. A general review of the KM algebra can be found in D. Gepner and E. Witten, *Nucl. Phys.* **B278**, 493 (1986); P. Goddard and D. Olive, *Inter. J. Mod. Phys.* **1**, 303 (1986); V.G. Kac, “Infinite dimensional Lie algebra”, (Birkhauser, Boston, 1983).
17. I. Affleck, *Phys. Rev. Lett.* **56**, 746 (1986).
18. I would like to thank E. Fradkin for pointing this out to me.